

## INDEX TO VOLUME 63

## AUTHORS

- Aasen, Kenneth: see Brown, Aasen and von Geldern Jr., 60-64
- Ainley, David G.: see Prerotti, Ainley and Lewis, 64-66
- Anderson, Daniel W., Ronald M. Jurek and James O. Keith: The Status of Brown Pelicans at Anacapa Island in 1975, 4-10
- Anderson, M. Eric: Range Extension of Two Marine Fishes to the Monterey Bay Area, 132-133
- Applegate, Shelton P.: A New Record-Size Bonito Shark, *Isurus oxyrinchus* Rafinesque, From Southern California, 126-129
- Bargmann, Gregory G.: Instances of Copper Rockfish Consuming a Spiny Dogfish Shark, 192
- Barnhart, Roger A.: see Rabin and Barnhart, 193-196
- Behrens, David W.: Fecundity and Reproduction of the Viviparous Perches *Hypsurus caryi* (Agassiz) and *Embiotoca jacksoni* Agassiz, 234-252
- Benville, Jr., Pete E., and Sid Korn: The Acute Toxicity of Six Monocyclic Aromatic Crude Oil Components to Striped Bass (*Morone saxatilis*) and Bay Shrimp (*Crango franciscorum*), 204-209
- Bertram, Ronald C., and Ronald D. Rempel: Migration of North Kings Deer Herd, 157-179
- Black, Glenn F.: Notes on Some Fishes Collected Off the Outer Coast of Baja California, 71-72
- Borneman, John C.: see Wilber, Mallette and Borneman, 189-190
- Brown, Delores, Kenneth Aasen and C. E. von Geldern Jr.: Alabama Spotted Bass Grow at Record Rate in Lake Perris, California, 60-64
- Brown, Delores, Edward E. Miller and C. E. von Geldern Jr.: Detection of Delayed Annulus Formation Among Bluegill, *Lepomis macrochirus*, Populations at Lake Nacimiento, California, 29-42
- Bryant, Gary Lee: Fecundity and Growth of Tule Perch, *Hysterothorax traski*, in the Lower Sacramento-San Joaquin Delta, 140-156
- Cailliet, Gregor M., and Robert N. Lea: Abundance of the "Rare" Zoarid, *Maynea californica* Gilbert, 1915, in the Monterey Canyon, Monterey Bay, California, 253-261
- DeMartini, John D.: see Giorgi and DeMartini, 80-94
- DeWitt, Leray A.: Notes on Shark Species in Nearshore Waters Near Point Conception, California, 273-274
- Duffy, John M.: Southern Range Extensions for Chum and Sockeye Salmon, *Oncorhynchus keta* and *O. nerka*, 196-199
- Eldridge, Maxwell B.: Factors Influencing Distribution of Fish Eggs and Larvae Over Eight 24-Hr Samplings in Richardson Bay, California, 101-116
- Floerke, Robert: Reported Cases of Avian Tuberculosis in California Waterfowl, 191-192
- Fry, Michael E., and Charles E. Vaughn: Acorn Selection by Band-Tailed Pigeons, 59-60
- Garcelon, David K.: An Expandable Drop-off Transmitter Collar for Young Mountain Lions, 185-189
- Giorgi, Albert E., and John D. DeMartini: A Study of the Reproductive Biology of the Red Abalone, *Haliotis rufescens* Swainson, Near Mendocino, California, 80-94
- Gotshall, Daniel W.: Stomach Contents of Northern California Dungeness Crabs, *Cancer magister*, 43-51
- Gray, Randall L.: Extension of Red Fox Distribution in California, 58
- Haaker, Peter L.: Abnormal Vertebral Development in Northern Anchovy, *Engraulis mordax* Girard, 182-185
- Hauser, William J.: Temperature Requirements of *Tilapia zillii*, 228-233
- Hennessy, Scott L., and G. Victor Morejohn: Acanthicephalan Parasites of the Sea Otter, *Enhydra lutris*, Off Coastal California, 268-272
- Hubbs, Carl L.: First Record of Mating of Ridley Turtles in California, with Notes on Commensals, Characters, and Systematics, 262-267
- Jurek, Ronald M.: see Anderson, Jurek and Keith, 4-10
- Keith, James O.: see Anderson, Jurek and Keith, 4-10
- Knudson, Peter M.: Observations on the Breeding of the Harbor Seal in Humboldt Bay, California, 66-70
- Korn, Sid: see Benville, Jr. and Korn, 204-209
- Kuljis, Andrew: see Lal, Lasker and Kuljis, 210-218
- Lal, Drishan, Reuben Lasker and Andrew Kuljis: Acclimation and Rearing of Striped Bass Larvae in Sea Water, 210-218
- Lasker, Reuben: see Lal, Lasker and Kuljis, 210-218
- Lea, Robert N.: see Cailliet and Lea, 253-267
- Lewis, T. James: see Prerotti, Ainley and Lewis, 64-66
- Mallette, Robert D.: see Wilber, Mallette and Borneman, 189-190

- Martin, John H.: see Martin, Stephenson and Martin, 95-100
- Martin, Michael, Mark D. Stephenson and John H. Martin: Copper Toxicity Experiments in Relation to Abalone Deaths Observed in a Power Plant's Cooling Waters, 95-100
- Mecum, W. Lee: see Meinz and Mecum, 277-278
- Meinz, Mike, and W. Lee Mecum: A Range Extension for Mississippi Silverside in California, 277-278
- Miller, Edward E.: see Brown, Miller and von Geldren Jr., 29-42
- Morejohn, G. Victor: see Hennessy and Morejohn, 268-272
- Oglesby, Larry C.: A Newly Introduced, Brackish-Water Snail in the Salton Sea Basin, California, 180-182
- Phelps, Robert: see Valentine and Phelps, 129-132
- Prerotti, Raymond J., David G. Ainley and T. James Lewis: Birth of a California Sea Lion on Southeast Farallon Island, 64-66
- Quelvog, Brian D.: First Record of the Striped Mullet (*Mugil cephalus*) in the Sacramento-San Joaquin Delta of California, 278-279
- Rabin, Douglas J., and Roger A. Barnhart: Fecundity of Pacific Herring, *Clupea harengus pallasi*, in Humboldt Bay, 193-196
- Rao Teegavarapu R.: Effects of Salinity on Larval Growth in the California Killifish, *Fundulus parvipinnis* Girard, 22-28.
- Rawstron, Robert R.: Effect of a Reduced Bay Limit and Later Planting Date on the Mortality, Survival, and Cost and Yield of Three Domestic Strains of Rainbow Trout at Lake Berryessa and Merle Collins Reservoir 1971-1974, 219-227; Harvest, Survival, and Weight Returns of Tagged Eagle Lake and Coleman Rainbow Trout Stocked in Lake Berryessa in 1972, 274-277
- Rempel, Ronald D.: see Bertram and Rempel, 157-179
- Reynolds, Linda J.: see Reynolds and Reynolds, 124-125
- Reynolds, William W., and Linda J. Reynolds: Observations on Food Habits of the Angelfishes, *Pomacanthus zonipectus* and *Holacanthus passer* in the Gulf of California, 124-125
- Shimek, Steven J.: The Underwater Foraging Habits of the Sea Otter, *Enhydra lutris*, 120-122
- Smith, Albert C.: Reactions of Fish Red Blood Cells with Mucus and Sera from Other Fish(es), 52-57
- Standord, Wilber R., Robert D. Mallette and John C. Borneman: California Condor Survey, 1975, 189-190
- Stephenson, Mark D.: Sea Otter Predation on Pismo Clams in Monterey Bay, 117-120; see Martin, Stephenson and Martin, 95-100
- Stienecker, Walter E.: Supplemental Data on the Food Habits of the Western Gray Squirrel, 11-21
- Valentine, David W., and Robert Phelps: Parasitism of Barred Surfperch, *Amphistichus argenteus*, By *Nerocila californica*: A Fit to the Poisson, 129-132
- Vaughn, Charles E.: see Fry and Vaughn, 59-60
- von Geldren Jr., C. E.: see Brown, Aasen and von Geldren Jr., 60-64; Brown, Miller and von Geldren Jr., 29-42
- Wicksten, Mary K.: Feeding in the Decorator Crab, *Loxorhynchus crispatus* (Brachyura: Majidae), 122-124

## SCIENTIFIC NAMES

- |   |   |
|---|---|
| <i>Abies concolor</i> : 13, 176                         | <i>Atherinops affinis</i> : 109               |
| <i>Abies magnifica</i> : 176                            | <i>Atherinopsis californiensis</i> : 109      |
| <i>Acanthochondria comuta</i> : 259                     | <i>Aythya affinis</i> : 191                   |
| <i>Accipiter cooperi</i> : 190                          | <i>Blepharipoda occidentalis</i> : 119        |
| <i>Accipiter striatus</i> : 190                         | <i>Branta canadensis</i> : 191                |
| <i>Aesculus californica</i> : 11                        | <i>Brevoortia tyrannus</i> : 113              |
| <i>Allosmerus elongatus</i> : 44, 45, 109               | <i>Brosomphycis marginata</i> : 109           |
| <i>Alpheus dentipes</i> : 122                           | <i>Buteo jamaicensis</i> : 190                |
| <i>Ambloplites rupestris</i> : 38                       | <i>Buteo regalis</i> : 190                    |
| <i>Ammodytes hexapterus</i> : 109                       | <i>Buteo swainsoni</i> : 190                  |
| <i>Amphistichus argenteus</i> : 129, 130, 247, 249, 250 | <i>Calocedrus decurrens</i> : 13              |
| <i>Anas acuta</i> : 191                                 | <i>Cancer magister</i> : 43, 45               |
| <i>Anas carolinensis</i> : 191                          | <i>Caretta caretta caretta</i> : 264          |
| <i>Anas cyanoptera</i> : 191                            | <i>Cathartes aura</i> : 190                   |
| <i>Anas platyrhynchos</i> : 191                         | <i>Ceanothus cordulatus</i> : 175             |
| <i>Aquila chrysaetos</i> : 190                          | <i>Ceanothus cuneatus</i> : 161               |
| <i>Arbacia punctulata</i> : 88                          | <i>Cephaloscyllium ventriosum</i> : 273-275   |
| <i>Arbutus menziesii</i> : 11                           | <i>Chamaebatia foliolosa</i> : 176            |
| <i>Archidoris montereyensis</i> : 123                   | <i>Chanos chanos</i> : 52, 53                 |
| <i>Arctocephalus tropicalis gazella</i> : 64            | <i>Chelonia agassizii</i> : 266               |
| <i>Artemia nauplii</i> : 22, 23                         | <i>Chelonia mydas</i> : 266                   |
| <i>Artemia salina</i> : 122                             | <i>Chelonia mydas carolinensis</i> : 265, 266 |
| <i>Astrometis sertulifera</i> : 124                     | <i>Chirolophis decoratus</i> : 139            |

- Chirolepis polyactocephalus*: 139  
*Circus cyaneus*: 190  
*Citharichthys sordidus*: 44, 45  
*Citharichthys stigmæus*: 44, 45  
*Clangula clangula*: 269  
*Clevelandia ios*: 113  
*Clinocardium nuttalli*: 54  
*Clupea harengus pallasii*: 101, 108, 109, 111, 112, 114, 115, 193  
*Collisella pelta*: 123  
*Columbia fasciata*: 59  
*Columbia palumbus*: 60  
*Conchoderma virgatum*: 265  
*Corynosoma enghdri*: 268–271  
*Coryphopterus nicholsi*: 132  
*Cottus asper*: 114  
*Crago franciscorum*: 204  
*Cylindrolepis darwini*: 265  
*Cymatogaster aggregata*: 109, 143, 146, 148, 246, 247, 249, 250  
*Cynoscion nobilis*: 108, 109, 113, 114  
*Cyprinodon macularius*: 27, 180  
*Cyprinus carpio*: 37  
*Damalichthys argyrosomus*: 143  
*Dendraster excentricus*: 123, 124  
*Diastylepis dawsoni*: 45  
*Distichlis spicata stricta*: 180  
*Ditrema temminckii*: 247, 249  
*Ditrema viridis*: 247  
*Dorosoma petenense*: 29, 210, 219, 274  
*Echinorhinus cookei*: 256  
*Egretta laevigata*: 260  
*Embiotoca jacksoni*: 234, 236, 241, 244–247, 249  
*Embiotoca lateralis*: 247, 249  
*Emerita analoga*: 271, 272  
*Engraulus mordax*: 44, 45, 101, 108, 109, 111, 113–115, 182–184  
*Enhydra lutris*: 117, 120, 268  
*Eptatretus stoutii*: 256  
*Eretmochelys imbricata*: 262  
*Eumetopias jubatus*: 64  
*Falco mexicanus*: 190  
*Falco sparverius*: 190  
*Filicollis anatis*: 271  
*Fulica americana*: 191  
*Fundulus parvipinnis*: 22, 27  
*Galeorhinus zyopterus*: 273–275  
*Gambusia affinis*: 180  
*Grus canadensis*: 191  
*Gymnogyps californianus*: 189  
*Haliotis cracherodii*: 86, 95  
*Haliotis cyclobates*: 88  
*Haliotis discus hannai*: 87  
*Haliotis gigantea*: 91  
*Haliotis iris*: 84, 87  
*Haliotis Kamtschatkana*: 84  
*Haliotis laevigata*: 88  
*Haliotis midas*: 87  
*Haliotis rufescens*: 80, 81, 95  
*Haliotis tuberculata*: 87, 91  
*Hemiphysa novaeseelandiae*: 60  
*Hennicia levisculia*: 123  
*Hexanchus griseus*: 256  
*Holacanthus bermudensis*: 125  
*Holacanthus ciliaris*: 125  
*Holacanthus clarionensis*: 124  
*Holacanthus passer*: 124, 125  
*Holacanthus tricolor*: 125  
*Hyas araneus*: 123  
*Hydrolagus collieri*: 256  
*Hyperprosopon argenteum*: 249  
*Hyperprosopon ellipticum*: 247  
*Hypospsetta guttulata*: 108–110, 112–114  
*Hypsurus caryi*: 234, 236, 240–247, 249, 250  
*Hysteroecarpus traski*: 140–156, 247, 249  
*Ictalurus punctatus*: 61  
*Ilypnus gilberti*: 110  
*Isurus belyaevi*: 128  
*Isurus oxyrinchus*: 126–129  
*Larus argentatus*: 269  
*Larus glaucescens*: 271  
*Larus occidentalis*: 64, 67, 271  
*Leiostomus xanthurus*: 113  
*Lepidochelys olivacea*: 262, 265, 266  
*Lepidogobius lepidus*: 114  
*Lepomis cyanellus*: 61  
*Lepomis macrochirus*: 29  
*Leptasterias aequalis*: 123  
*Leptocottus armatus*: 108, 110, 113, 114  
*Logodon rhomboides*: 113  
*Loxorhynchus crispatus*: 122–124  
*Loxorhynchus grandis*: 123  
*Lutianus fulvus*: 52, 53  
*Lyopsetta exilis*: 110, 114  
*Lythrypnus dalli*: 133  
*Lythrypnus zebra*: 132, 133  
*Macoma balthica*: 88  
*Mareca americana*: 191  
*Macrocypris pyriferus*: 260  
*Maynea bulbiceps*: 256  
*Maynea californica*: 253, 255–261  
*Melanitta deglandi*: 269  
*Melanitta perspicillata*: 269, 271  
*Melanostigma pammelas*: 258  
*Menidia audens*: 277  
*Microgadus proximus*: 44, 45  
*Micrometrus aurora*: 247  
*Micrometrus minimus*: 247  
*Micropterus dolomieu*: 38, 60  
*Micropterus punctulatus*: 38  
*Micropterus punctulatus henshalli*: 60  
*Micropterus punctulatus punctulatus*: 60  
*Micropterus salmoides*: 29  
*Mimulus foliatus*: 123  
*Miroganga angustirostris*: 64  
*Mitrella tuberosa*: 45  
*Morone chrysops*: 29, 210  
*Morone saxatilis*: 101, 204, 210  
*Mugil cephalus*: 52, 278  
*Mycobacterium tuberculosis*: 191  
*Mytilus californicus*: 7  
*Mytilus edulis*: 122  
*Nerocila californica*: 129–132  
*Oligocottus maculosus*: 110  
*Oligocottus snyderi*: 110  
*Olivella biplicata*: 121  
*Oncorhynchus keta*: 196  
*Oncorhynchus kisutch*: 27, 199  
*Oncorhynchus nerka*: 198  
*Ophiodon elongatus*: 110, 191  
*Ondatra zibethica*: 180  
*Orchomene obtusa*: 259  
*Oxyjulis californica*: 71  
*Oxyura jamaicensis*: 191  
*Paralichthys californicus*: 110, 114  
*Parophrys vetulus*: 110, 114  
*Pelagia noctiluca*: 123  
*Pelecus occidentalis californicus*: 4  
*Peprilus similis*: 110, 114  
*Rhacochilus vacca*: 247  
*Phanerodon furcatus*: 247, 250  
*Phoca vitulina richardi*: 66, 68  
*Phoca vitulina vitulina*: 69  
*Phyllospadix scouleri*: 253, 256  
*Phyllospadix torreyi*: 260  
*Pinus lambertiana*: 13  
*Pinus ponderosa*: 13  
*Pinus sabiniana*: 13  
*Planes cyaneus*: 264  
*Planes minutus*: 265  
*Platichthys stellatus*: 110, 114  
*Pleurobranchaea californica*: 259  
*Poecilia sphenops*: 27  
*Polymorphus altmani*: 269–272

- Polymorphus kenti*: 269-272  
*Polymorphus major*: 269  
*Pomacanthus arcuatus*: 125  
*Pomacanthus paru*: 125  
*Pomacanthus zonipectus*: 124, 125  
*Pomoxis annularis*: 38  
*Pomoxis nigromaculatus*: 29, 226  
*Porichthys notatus*: 109  
*Prionace glauca*: 274  
*Prionotus gymnotesthus*: 71  
*Prunus emarginata*: 176  
*Psettichthys melanostictus*: 110, 113, 114  
*Pugettia producta*: 123  
*Pycnopodia helianthoides*: 123  
*Quercus agrifolia*: 11  
*Quercus douglasii*: 11, 59  
*Quercus kelloggii*: 11, 60, 176  
*Quercus lobata*: 11  
*Quercus vaccinifolia*: 176  
*Quercus wislizenii*: 59  
*Raja inornata*: 71  
*Remora remora*: 264  
*Rhacochilus vacca*: 243, 246, 249  
*Salmo gairdneri*: 61, 219, 274  
*Salmo gairdneri aquilarum*: 274  
*Sciurus griseus*: 11  
*Scorpaenichthys marmoratus*: 110, 112, 122  
*Sebastes caurinus*: 192  
*Sebastes rosenae*: 132  
*Siliqua patula*: 45  
*Spatula clypeata*: 191  
*Spirinchus starksi*: 44, 45  
*Spirinchus thaleichthys*: 109  
*Spisula solidissima*: 84  
*Squalus acanthias*: 192, 273, 275  
*Squatina californica*: 273-275  
*Strongylocentrotus franciscianus*: 123, 124  
*Strongylocentrotus purpuratus*: 122, 123  
*Syngnathus leptorhynchus*: 109, 113  
*Tachysurus australis*: 55  
*Thaleichthys pacificus*: 133  
*Thiara granifera*: 180, 181, 182  
*Tilapia mossambica*: 52, 53  
*Tilapia zillii*: 180, 228-232  
*Tivela stultorum*: 117  
*Tresus capax*: 84  
*Triakis semifasciata*: 273-275  
*Tropia maculata*: 123  
*Typha domingensis*: 180  
*Umbellularia californica*: 11  
*Urechis caupo*: 121  
*Veilella veilella*: 124  
*Vulpes fulva*: 58  
*Vulpes fulva necator*: 58  
*Yoldia limatula*: 45  
*Zalophus californianus*: 64  
*Zaniclepis latipinnis*: 71  
*Zostera marina*: 45

## SUBJECT

- Abalone, red: study of the reproductive biology near Mendocino, California, 80-94  
 Acclimation: and rearing of striped bass larvae in sea water, 210-218  
 Accorn: selection by band-tailed pigeons, 59-60  
 Anacapa Island: status of brown pelicans at, in 1975, 4-10  
 Anchovy, northern: abnormal vertebral development 182-185  
 Angelfishes: in the gulf of California, food habits of, 124, 125  
 Annulus formation: delayed among bluegill populations at Lake Nacimiento, California, 29-42  
 Avian tuberculosis: reported cases in California waterfowl, 191-192  
 Baja California: notes on some fishes collected off the outer coast of, 71-72  
 Bass, Alabama spotted: grow at record rate in Lake Perris, California, 60-64  
 Bass, striped: and bay shrimp acute toxicity of six monocyclic aromatic oil components to, 204-209; larvae acclimation and rearing in sea water, 210-218  
 Bluegill: delayed annulus formation among, populations at Lake Nacimiento, 29-42  
 Breeding behavior: of the harbor seal in Humboldt Bay, 66-70  
 Condor, California: survey, 1975, 189-190  
 Cost: mortality, survival, and yield of three domestic strains of rainbow trout at Lake Berryessa and Merle Collins Reservoir 1971-1974, 219-227  
 Crab, decorator: feeding in the, 122-124  
 Crab, northern California dungeness: stomach contents of, 43-52  
 Farallon Island, southeast: birth of a California sea lion, 64-66  
 Fecundity: of tule perch in the lower Sacramento-San Joaquin Delta, 140-156; of Pacific herring in Humboldt Bay, 193-196; and reproduction of the viviparous perches, 234-252  
 Feeding: in the decorator crab, 122-124  
 Fish eggs and larvae: over eight 24-hr samplings in Richards Bay, California, 101-116  
 Fishes collected: off the outer coast of Baja California, notes on, 71-72  
 Food habits: of the Western gray squirrel, 11-21; of the angelfishes in the gulf of California, 124-125  
 Foraging habits: of the sea otter underwater, 120-122  
 Gray squirrel, Western: supplemental data on food habits of, 11-21  
 Growth: of tule perch in the lower Sacramento-San Joaquin Delta, 140-156  
 Harvest: survival and weight returns of tagged Eagle Lake and Coleman rainbow trout stocked in Lake Berryessa in 1972, 274-277

- Herring, Pacific: fecundity of, in Humboldt Bay, 193-196
- Humboldt Bay: breeding behavior of the harbor seal in, 66-70; fecundity of Pacific herring, 193-196
- Killifish: effects of salinity on larval growth in California, 22-29
- Lake Nacimiento: delayed annulus formation among Bluegill populations at, 29-42
- Lake Perris: Alabama spotted bass grow a record rate in, 60-64
- Larval: growth in California killifish, 22-29
- Mating: of ridley turtles in California, first record of, 262-267
- Mendocino, California: a study of the reproductive biology of the red abalone, 80-94
- Migration: of North kings deer herd, 157-179
- Mississippi silversides: range extension in California, 277-278
- Monocyclic aromatic crude oil components: to striped bass and bay shrimp, the acute toxicity of, 204-209
- Monterey Bay: sea otter predation on pismo clams in, 117-120; range extension of two marine fishes to the, 132-133
- Monterey Canyon: abundance of "rare" zoarid in, 253-261
- Mortality: survival, cost and yield of three domestic strains of rainbow trout at Lake Berryessa and Merle Collins Reservoir 1971-1974, 219-227
- Mountain lions, young: an expandable drop-off transmitter collar for, 185-189
- Mullet, striped: first record of, in Sacramento-San Joaquin Delta of California, 278-279
- North kings deer: herd migration of, 157-179
- Parasites, acanthocephalan: of sea otter off coastal California, 268-272
- Parasitism: of barred surfperch by *Nerocila californica*: a fit to the poisson, 129-132
- Pelicans, brown: at Anacapa Island 1975, status of, 4-10
- Perch, tule: fecundity and growth in the lower Sacramento-San Joaquin Delta, 140-156
- Perches, viviparous: fecundity and reproduction of, 234-252
- Pigeons, band-tailed: acorn selection by, 59-60
- Pismo clams: in Monterey Bay, sea otter predation on, 117-120
- Point Conception: shark species in nearshore waters near, 273-274
- Range extension: of two marine fishes to the Monterey Bay area, 132-133; for Mississippi silverside in California, 277-278
- Range extensions, southern: for chum and sockeye salmon, 196-199
- Rearing: and acclimation of striped bass larvae in sea water, 210-218
- Red blood cells, fish: reactions of, with mucus and sera from other fish(es), 52-57
- Red Fox: distribution in California, extension of, 58
- Reproduction: biology of the red abalone near Mendocino, California, 80-94; and fecundity of viviparous perches, 234-252
- Reviews: Mammals of the world, 73; Shad fishing, 73; The fishes of Missouri, 73; The soft-hackled fly, 74; The practical fly fisherman, 74; Culture of marine invertebrate animals, 74; The ageing of fish: proceedings of an international symposium, 75; Marine game fishes of the Pacific coast from Alaska to the equator, 75; Fishes in Kansas, 76; A field guide to the common and interesting plants of Baja California, 134; Modern fly dressings for the practical angler, 134; Fishing with light, 134; The ageing of fish, 200; Marine game fishes of the Pacific coast from Alaska to the equator, 200; Coastal fishes of Southern Japan, 280; Fishes of the world, 281; Waterfowl of North America, 281; Advanced Bass Fishing, 281; Fly tackle, 282; Classic rods and rodmakers, 282; The essential fly tier, 282
- Richards Bay, California: fish eggs and larvae over eight 24-hr samplings in, 101-116
- Rockfish, copper: instances of consuming a spiny dogfish shark, 192
- Sacramento-San Joaquin Delta: fecundity and growth of tule perch, 140-156; first record of the striped mullet in, 278-279
- Salinity: on larval growth in California killifish, 22-29
- Salmon, chum and sockeye: southern range extensions, 196-199
- Salton Sea Basin, California: a newly introduced brackish-water snail, 180-182
- Samplings: of fish eggs and larvae in Richards Bay, California, 101-116
- Seal, harbor: breeding behavior of, in Humboldt Bay, California, 66-70
- Sea Lion, California: birth of, on Southeast Farallon Island, 64-66
- Sea otter: predation on pismo clams in Monterey Bay, 117-120; underwater foraging habits of, 120-122; acanthocephalan parasites of, off coastal California, 268-272

- Shark: species in nearshore waters near Point Conception, 273-274
- Shark, bonito: new record size from Southern California, 126-129
- Shark, spiny dogfish: instances of copper rockfish consuming, 192
- Shrimp, bay: and striped bass acute toxicity of six monocyclic aromatic crude oil components to, 204-209
- Snail, brackish-water: newly introduced in the Salton Sea Basin, California, 180-182
- Status: of brown pelicans at Anacapa Island in 1975, 4-10
- Stomach contents: of Northern California dungeness crabs, 43-52
- Surfperch, barred: parasitism by *Nerocila californica*: a fit to poisson, 129-132
- Survey: California condor, 1975, 189-190
- Survival: mortality, cost and yield of three domestic strains of rainbow trout at Lake Berryessa and Merle Collins Reservoir 1971-1974, 219-227; harvest and weight returns of tagged Eagle Lake and coleman rainbow trout stocked in Lake Berryessa in 1972, 274-277
- Temperature: requirements of *Tilapia zillii*, 228-233
- Tilapia zillii*: temperature requirements of, 228-233
- Toxicity: of six monocyclic aromatic crude oil components to striped bass and bay shrimp, 204-209
- Toxicity, copper: experiments in abalone deaths observed in power plant's cooling waters, 95-100
- Transmitter collar, expandable drop-off: for young mountain lions, 185-189
- Trout, rainbow: mortality, survival, cost and yield of three domestic strains at Lake Berryessa and Merle Collins Reservoir 1971-1974, 219-227; harvest, survival, and weight returns of tagged Eagle Lake and coleman stocked in Lake Berryessa in 1972, 274-277
- Tuberculosis, avian: reported cases in California waterfowl, 191-192
- Turtles, ridley: first record of mating in California, 262-267
- Vertebral: abnormal development in Northern anchovy, 182-185
- Warbonnet, decorated: ERRATUM for currently accepted AFS/ASH name is *Chirolophis polyactcephalus*, 139
- Waterfowl, California: reported cases of avian tuberculosis, 191-192
- Weight returns: of tagged Eagle Lake and coleman rainbow trout in Lake Berryessa in 1972, 274-277
- Yield: mortality, survival, and cost of three domestic strains of rainbow trout at Lake Berryessa and Merle Collins Reservoir 1971-1974, 219-227
- Zooid: abundance of, in Monterey Canyon, 253-261

